

CANIQUEL[®] Plus XL tablets ad us. vet.

QUALITATIVE AND QUANTITATIVE COMPOSITION

One tablet contains

Praziquantel	100.0 mg
Fenbendazole	1000.0 mg

Pharmaceutical form

Tablets for oral use



CLINICAL PARTICULARS

Target species

Dogs

Indications of use

For the treatment of diagnosed mixed infections with roundworms and intestinal stages of tapeworms in dogs

<u>Ascarids</u>	<i>Toxocara canis</i> <i>Toxascaris leonina</i>
<u>Hookworms</u>	<i>Ancylostoma caninum</i> , <i>Uncinaria stenocephala</i>
<u>Whipworms</u>	<i>Trichuris vulpis</i>
<u>Tapeworms</u>	<i>Echinococcus</i> spp. <i>Dipylidium caninum</i> <i>Taenia</i> spp. <i>Mesocestoides</i> spp.

Contraindications

Due to the lack of relevant informations Caniquantel[®] Plus XL tablets should not be used in pregnant animals or in pups under three weeks of age.

Posology and route of administration

The recommended dosage is 5 mg of Praziquantel and 50 mg of Fenbendazole per kg body weight. This corresponds to 1 tablet Caniquantel plus XL per 20 kg body weight or a half tablet for 10 kg bodyweight.

The tablet is administered directly or mixed with food. Dietary measures or fasting are not necessary. For a treatment of infected dogs a repeated administration on three consecutive days is recommended.

Use during pregnancy and lactation

Not to be used in pregnant animals

Withdrawal period(s)

Not applicable

PHARMACEUTICAL PARTICULARS

Shelf life

Shelf-life of the veterinary medicinal product as packaged for sale: 3 years

Special precautions for storage

Store at room temperature (15 – 25°C)

Protected from light.

Presentations

1 Blister containing 3 tablets

(The presentation varies from country to country)